

A Study of *Sarcocystis* Infection in Mincemeat Using Digestion Method in Ghazvin, Iran

Jaber Davoudi^a, Afshin Bahman Shabestari^a, Peyman Mozahheb^a, Roghayeh Norouzi^b,
Mohammadreza Maadinia^a, Razzagh Mahmoudi^{c*}

^aDepartment of Veterinary Parasitology, Islamic Azad University, Abhar-Branch, Abhar, Iran.

^bDepartment of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran.

^cMedical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

*Correspondence should be addressed to Dr. Razzagh Mahmoudi, Email: r.mahmodi@yahoo.com

A-R-T-I-C-L-E-I-N-F-O

Article Notes:

Received: Sep. 26, 2016

Received in revised form:
Jan. 16, 2017

Accepted: Feb. 24, 2017

Available Online: Feb 28,
2017

Keywords:

Sarcocystis infection
Mincemeat
Digestion method
Ghazvin
Iran

A-B-S-T-R-A-C-T

Background & Aims of the Study: Sarcocystiosis is a zoonosis appeared in domestic animals caused by various species of *Sarcocystis*. This protozoan disease has worldwide distribution among human and many species of animals. Humans acquire infection by eating of raw and under cooked beef, pork or mincemeat containing schizonts of *Sarcocystis hominis* and *S. suihominis*. The aim of present study is to detect prevalence of the *Sarcocystis* spp. infection in mincemeat samples at Ghazvin province of Iran.

Materials & Methods: Three hundred mincemeat samples of 150 sheep and 150 cattle were collected from butchers (in spring 2013) in different areas of Ghazvin province, Iran. The statistical analysis was done by independence sample t test, using SPSS ver. 22.0.0 (Chicago, IL, USA).

Results: the finding of this study showed that the highest prevalence of *Sarcocystis* infection rate was observed in cattle (92.8%) and the lowest of that was evident in sheep (85.6%). The highest infection rates in both types of minced meat samples were in May (45 and 49 minced meat of sheep and cattle, respectively).

Conclusions: The results revealed that Ghazvin province has the highest *Sarcocystis* infection rate. Regarding to the high prevalence of *Sarcocystis* contamination in this study, prevention of eating raw or under-cooked meat is strongly recommended.

Please cite this article as: Davoudi J, Bahman Shabestari A, Mozahheb P, Norouzi R, Maadinia M, Mahmoudi R. A study of sarcocystis infection in mincemeat using digestion method in Ghazvin, Iran. Arch Hyg Sci 2017;6(2):178-181.